

Are there any complications ?

As with any other minimally invasive procedures, side effects of Sacroiliac Joint Neurotomy treatment using the Pain Management SInergy™ System are relatively rare. They could include infection, inflammation, and tenderness at the treatment site. Side effects are typically mild, and most patients generally feel better within a few days after the procedure. If you experience any side effects such as severe pain at the site of needle insertion, severe headaches, or swelling you should contact your doctor immediately.



When can I return to normal activity ?

Following consultation with your doctor, a return to normal activity may be possible one or two days after the procedure. It may be recommended that you pursue a course of physical therapy and avoid certain strenuous activities.

Baylis Medical is a world leader in the development and distribution of high-technology pain management products. We conceive, innovate, develop, manufacture, and sell state-of-the-art medical products used around the world. Baylis Medical operates under the ISO 13485:2003 approved quality system.

Baylis
MEDICAL

Baylis Medical Company Inc.
5959 Trans-Canada Highway
Montreal, QC
Canada
H4T 1A1

Tel : (514) 488-9801
Fax : (514) 488-7209
www.baylismedical.com
info@baylismedical.com


Find more information, visit our Web site
www.sinergysystem.com

The BMC logo, SInergy™ and SInergy™ System are trademarks or registered trademarks of Baylis Medical Company Inc. in the United States and/or other countries.

© Copyright Baylis Medical Company Inc., 2008
Patents Pending and/or issued,


May be a solution for your pain

SACROILIAC JOINT RF NEUROTOMY



Description of sacroiliac joint pain

Chronic sacroiliac joint pain can be a frustrating and debilitating condition that is often hard to diagnose. Symptoms of sacroiliac joint (SIJ) pain are varied, and can include a severe aching discomfort across the lower back and buttock region that is usually worse



in the early morning and after extended periods of sitting. Rest does not usually relieve chronic SIJ pain, and this dull, constant pain can radiate throughout the entire lower back region.

Some estimates are that sacroiliac joint pain affects up to 30% of lower back pain sufferers. For some patients the pain subsides after a few months or with conservative treatment such as bed rest, or anti-inflammatory medication. For others the pain becomes chronic (lasting longer than six months) and may necessitate more invasive treatments. Sacroiliac Joint RF Neurotomy Treatment using the Pain Management SInergy™ System is a new and alternative solution to manage chronic sacroiliac joint pain.

What is RF neurotomy for SIJ pain ?

RF neurotomy uses radiofrequency energy to safely and effectively disable the sensory nerves that may be responsible for SIJ pain. During the procedure, the patient lies on a table and usually mildly sedated. The area to be treated is numbed and SInergy™ Probes, developed by Baylis Medical, are placed into the back through an introducer needle. The painful nerves are then treated with gentle warming by radiofrequency waves. After the treatment, the probe and needles are removed and a bandage is placed over the treatment site. Disabling the sensory nerves in your SIJ region can potentially reduce your pain, and restore your ability to lead a normal life.

How long does the procedure take ?

It takes approximately 45 minutes to treat a painful sacroiliac joint. It is performed as an outpatient procedure, does not involve general anesthesia, and patients normally return home shortly after the procedure.

Find more information, visit our Web site
www.sinergysystem.com

Who can benefit from this treatment ?

If you have chronic sacroiliac joint pain, you may benefit from Sacroiliac Joint Neurotomy treatment using the Pain Management SInergy™ System. However, only a qualified physician can properly diagnose you and suggest appropriate treatments. Talk to your pain management physician to see if RF treatment for SIJ pain is right for you.

What do I do after the procedure ?

Because this procedure is minimally invasive, recovery is usually quick for most patients. However, you may be groggy from the sedative used during the procedure; and therefore require a responsible adult to drive you home afterwards. You may not drive or operate a motor vehicle on the day of your procedure.

Following the procedure, patients will normally be sore at the treatment site. This is to be expected because the insertion of the needles may cause inflammation, and the heat from the treatment may cause some short-term tenderness. The use of cold packs and analgesics may be suggested by your physician. Treatment site discomfort should subside within 2 to 7 days, but may last as long as two weeks. Patients typically experience some level of relief from their SIJ pain within 2 to 10 days of the procedure.